

碩士學位請求論文

ON THE ARC LENGTH UNDER INVERSION

指導教授 玄 進 五



濟州大學校 教育大學院

數學教育專攻

梁 昌 洪

1993年 8月

ON THE ARC LENGTH UNDER INVERSION

指導教授 玄 進 五

이 論文을 教育學 碩士學位 論文으로 提出함

1993年 6月 日


濟州大學校 教育大學院 數學教育專攻

提出者 梁 昌 洪




梁昌洪의 教育學 碩士學位 論文을 認准함

1993年 7月 日

審査委員長 양 성 호 

審査委員 김 현 수 

審査委員 玄 進 五 

CONTENTS

<Abstract>

Introduction	1
1. The arc length of a regular curve	2
2. The properties of inverse curve under inversion	6
3. The arc length under inversion	11
REFERENCES	15
< 초 목 >	16



<Abstract>

ON THE ARC LENGTH UNDER INVERSION

Yang, Chang-Hong

Mathematics Education Major
Graduate School of Education, Cheju National University
Cheju, Korea

Supervised by professor Hyen, Jin-Oh

Two points P and P' of the plane are said to be inverse with respect to a given circle $(O)_R$, if $OP \cdot OP' = R^2$ and also if both points are on the same side of O . Circle $(O)_R$ is called the circle of inversion and the transformation which sends point P into point P' is known as an inversion.

In this paper we consider the curves in two dimensional Euclidean space R^2 and prove that the length of a regular new curve segment $\beta(t)$ of the inside curve $\alpha(t)$ under inversion is equal to the length of a regular curve segment $\alpha(t)$ by scalar multiple.

Introduction

In this paper, our study of curves will be restricted to the certain plane curves in two dimensional Euclidean space R^2 .

In Section 1, we present the basic definitions and examples with respect to reparametrized curves and study some properties of the differential geometry, in particular, the arc length of curve segment

$$\alpha : [a, b] \rightarrow R^2.$$

Next, in Section 2, we introduce the definition and some properties of inverse curve under inversion. That is, the symbol $(O)_R$ is an inversion circle with center O and radius R and the relation between the inside point P and the outside point P' of $(O)_R$ is given by $OP \cdot OP' = R^2$ where its two points and O are collinear.

Finally, in Section 3, from the definition and the properties in Section 2, we prove the main theorem ; the length of a regular new curve segment $\beta(t)$ of the inside curve $\alpha(t)$ under inversion is equal to the length of a regular curve segment $\alpha(t)$ by scalar multiple.

1. The arc length of a regular curve

Let α be an injective function from an interval into R^2 and $\alpha(t)$ denote the curve in the plane. Then we have the derivative $\frac{d\alpha}{dt}(t_0)$ of α evaluated at $t = t_0$ if $\alpha(t)$ is differentiable in interval (a, b) .

Definition 1.1 A curve $\alpha : (a, b) \rightarrow R^2$ is called a regular curve if $\alpha \in C^k$ for some $k \geq 1$ and if $\frac{d\alpha}{dt} \neq 0$ for all $t \in (a, b)$.

If t is time, then the velocity vector of a regular curve $\alpha(t)$ at $t = t_0$ is the derivative evaluated at $t = t_0$. The speed of $\alpha(t)$ at $t = t_0$ is the length of the velocity vector at $t = t_0$, $\left| \frac{d\alpha}{dt}(t_0) \right|$.

Let $g : (c, d) \rightarrow (a, b)$ be an one-to-one and onto function, and let g and its inverse $h : (a, b) \rightarrow (c, d)$ be of class C^k for some $k \geq 1$. Then g is called a reparametrization of a curve $\alpha : (a, b) \rightarrow R^2$.

Proposition 1.2 If $\alpha : (a, b) \rightarrow R^2$ is a regular curve then the new curve $\beta = \alpha \circ g$ is a regular curve, if $\frac{dg}{dr} \neq 0$.

Proof.

$$(1.1) \quad \frac{d\beta}{dr} = \frac{d}{dr}[\alpha \circ g(r)] = \frac{d\alpha}{dt} \cdot \frac{dg}{dr},$$

that is,

$$\text{if } \frac{dg}{dr} \neq 0 \text{ then } \frac{d\beta}{dr} \neq 0.$$

Example 1.3 Let $g : (0, 1) \rightarrow (1, 2)$ be given by

$g(r) = 1 + r^2$. Then g is one-to-one and onto with inverse

$h(t) = \sqrt{t-1}$, $g \in C^k$ on $(0, 1)$ and $h \in C^k$ on $(1, 2)$ for some $k \geq 1$.

Thus g is a reparametrization of any regular curve on $(1, 2)$.

A regular curve segment is a function $\alpha : [a, b] \rightarrow R^2$ together with an open interval (c, d) , with $c < a < b < d$, and a regular curve $r : (c, d) \rightarrow R^2$ such that $\alpha(t) = r(t)$ for all $t \in [a, b]$.

Definition 1.4 The length of a regular curve segment $\alpha : [a, b] \rightarrow$

R^2 is defined by

(1.2)  제주대학교 중앙도서관
JEJU NATIONAL UNIVERSITY LIBRARY

$$\int_a^b \left| \frac{d\alpha(t)}{dt} \right| dt.$$

Theorem 1.5. The length of a curve is a geometric property, that is, it does not depend on the choice of reparametrization.

Proof. Let $g : [c, d] \rightarrow [a, b]$ be a reparametrization of a curve segment $\alpha : [a, b] \rightarrow R^3$, and let the new curve $\beta = \alpha \circ g$. Then, for $r \in [c, d]$,

since $g(r) = t$, $t \in [a, b]$, the length of β is

$$\begin{aligned}\int_c^d \left| \frac{d\beta}{dr} \right| dr &= \int_c^d \left| \frac{d}{dr}(\alpha \circ g) \right| dr \\ &= \int_c^d \left| \left(\frac{d\alpha}{dt} \right) \left(\frac{dg}{dr} \right) \right| dr \\ &= \int_c^d \left| \frac{d\alpha}{dt} \right| \left| \frac{dg}{dr} \right| dr.\end{aligned}$$

If $\frac{dg}{dr} > 0$, then $\left| \frac{dg}{dr} \right| = \frac{dg}{dr}$ and

$$g(c) = a, \quad g(d) = b.$$

Thus

$$\begin{aligned}\int_c^d \left| \frac{d\alpha}{dt} \right| \left| \frac{dg}{dr} \right| dr &= \int_c^d \left| \frac{d\alpha}{dt} \right| \left(\frac{dg}{dr} \right) dr \\ &= \int_a^b \left| \frac{d\alpha}{dt} \right| dt.\end{aligned}$$

If $\frac{dg}{dr} < 0$, then $\left| \frac{dg}{dr} \right| = -\frac{dg}{dr}$ and

$$g(c) = b, \quad g(d) = a.$$

Hence

$$\begin{aligned}\int_c^d \left| \frac{d\alpha}{dt} \right| \left| \frac{dg}{dr} \right| dr &= - \int_b^a \left| \frac{d\alpha}{dt} \right| \left(\frac{dg}{dr} \right) dr \\ &= \int_a^b \left| \frac{d\alpha}{dt} \right| dt.\end{aligned}$$

Example 1.6. Let $\alpha(t) = (r \cos t, r \sin t)$ with $r > 0$. Then $\frac{d\alpha}{dt} = (-r \sin t, r \cos t)$. Consider the arc length $s = s(t)$ of $\alpha(t)$.

Then

$$\begin{aligned}s &= \int_c \left| \frac{d\alpha}{dt} \right| dt \\ &= \int_c \sqrt{r^2 \sin^2 t + r^2 \cos^2 t} dt \\ &= rt.\end{aligned}$$

That is,



$$s = rt \text{ and } t = g(s) = \frac{s}{r}.$$

Hence,

$\beta(s) = (r \cos \frac{s}{r}, r \sin \frac{s}{r})$ is the unit speed parametrization of a circle of radius r .

2. The properties of inverse curve under inversion

In order to study the theorems in Section 3, we will see the properties of inverse curve.

Let the symbol $(O)_R$ denote the circle with center O and radius R .

Definition 2.1 Two points P and P' of the plane are said to be inverse with respect to a given circle $(O)_R$, if $OP \cdot OP' = R^2$ and if P, P' are on the same side of O and the (O, P, P') are collinear.

A circle $(O)_R$ is called the circle of inversion, and the transformation which sends point P into P' is called an inversion. As point P moves on a curve C , its inverse point P' moves on a curve C' which is the inverse curve of C . But the center O of the circle of inversion has no inverse point, for when P is at point O , $OP = 0$ and the relation $OP' = \frac{R^2}{OP}$ is meaningless.

Proposition 2.2 A line through O inverts into a line through O .

Proof. It is evident from the fact that O and inverse points are collinear.

Proposition 2.3 A line not through O inverts into a circle through O . Conversely, a circle through O inverts into a line not through O .

Proof. Let l be a line not through O and Q be the foot of the perpendicular from O to l , and let P be any point on l (Fig. 2.1).

Then, there are the inverse point Q' and P' of Q and P , respectively.

That is,

$$(2.1.a) \quad OQ \cdot OQ' = OP \cdot OP' = R^2$$

and

$$(2.1.b) \quad \frac{OQ}{OP} = \frac{OP'}{OQ'}$$

Therefore, $\triangle OQP$ and $\triangle OQ'P'$ have a common angle $\angle POQ$.


By (2.1), $\triangle OQP$ is similar to $\triangle OP'Q'$.

Thus

$$\angle OQP = \angle OP'Q' = 90^\circ.$$

But the arc in which a 90° angle is inscribed is a semicircle. Thus the point P' lies on a circle whose diameter is OQ' .

A reversal of these arguments completes the proof of this theorem.



< Fig. 2.1 >

Proposition 2.4 The angle between any two curves intersecting at a point which is different from the center O of the circle of inversion is unchanged under inversion.

Proof. Let the given curves C_1 and C_2 (Fig.2.2) intersect in a point P distinct from the center O of the circle of inversion and let any line l through O intersect these curves in the respective points A and B . Then the inverse curves to C_1 and C_2 , namely C'_1 and C'_2 , intersect at the inverse point P' to P .

If curves C'_1 and C'_2 are met by line l in the inverse points A' and B' of A and B , respectively. Let θ be the angle between the tangents

at P to curves C_1 and C_2 and let θ' be the angle between the tangents at P' to curves C'_1 and C'_2 . We must show that $\theta = \theta'$. Consider the triangles OPA and $OP'A'$. Then we have

$$(2.2) \quad \frac{OA}{OP} = \frac{OP'}{OA'}$$

Hence $\triangle OPA$ and $\triangle OP'A'$ are similar, so are $\triangle OPB$ and $\triangle OP'B'$.

Therefore

$$(2.3) \quad \angle OPA = \angle OA'P'$$


and

$$(2.4) \quad \angle OPB = \angle OB'P'$$

Subtraction (2.3) from (2.4) gives

$$\angle APB = \angle A'P'B'$$

Therefore $\lim_{l \rightarrow OP} \angle APB = \theta$ and $\lim_{l \rightarrow O'P'} \angle A'P'B' = \theta'$. Hence the proof is complete.



< Fig. 2.2. >

3. The arc length under inversion

Let $\alpha : (a, b) \rightarrow \mathbb{R}^2$ be the curve C_1 inside of inversion circle $(O)_R$. Then, for all $t \in (a, b)$, $\alpha(t)$ the image of α is the points P_t on curve C_1 . There exists an inverse curve $C_2 = \beta(t)$ outside of $(O)_R$.

Let OP_t be a distance from O to point P_t on curve C_1 . If a function $g : C_1 \rightarrow C_2$ is defined by

$$(3.1) \quad g(P_t) = P'_t \text{ for } P_t \in C_1,$$

then we can take a new curve $\beta(t) = g \circ \alpha(t)$ and see that the following properties hold.

Theorem 3.1 If curve $C_1 = \alpha(t)$ is a regular curve, then the inverse curve $C_2 = \beta(t)$ is also a regular curve.

Proof. Let $\alpha(t) = P_t$ for each $t \in (a, b)$. Then $\frac{d\alpha(t)}{dt} \neq 0$ for all $t \in (a, b)$, since $\alpha(t)$ is regular on (a, b) . Since $g(x, y) = \left(\frac{R^2 x}{x^2 + y^2}, \frac{R^2 y}{x^2 + y^2} \right)$, g is of class C^1 in $\mathbb{R}^2 \setminus \{(0, 0)\}$.

Now

$$\frac{d\beta(t)}{dt} = \begin{pmatrix} \frac{R^2(y^2 - x^2)}{(x^2 + y^2)^2} & \frac{-2R^2xy}{(x^2 + y^2)^2} \\ \frac{-2R^2xy}{(x^2 + y^2)^2} & \frac{R^2(x^2 - y^2)}{(x^2 + y^2)^2} \end{pmatrix} \frac{d\alpha(t)}{dt}.$$

Since $\frac{R^4(y^2 - x^2)(x^2 - y^2)}{(x^2 + y^2)^4} - \frac{4R^4x^2y^2}{(x^2 + y^2)^4} \neq 0$ for all (x, y) except

$(x, y) = (0, 0)$, $\frac{d\beta}{dt} \neq (0, 0)$, and hence β is regular in (a, b) .

Let OP_t and OP'_t be distances from the center of inversion circle $(O)_R$ to point P_t and P'_t on curves C_1 and C_2 , respectively. Consider the curve equation $OP_t = \alpha(t)$ with respect to the polar coordinate. Then the equation of the new curve $\beta(t)$ is given by $OP'_t = \beta(t)$.

Theorem 3.2 The length of a regular curve segment of new curve $\beta(t)$ of the inside curve $\alpha(t)$ under inversion is given by

$$\int_{t_1}^{t_2} \left| \frac{d\beta(t)}{dt} \right| dt = R^2 \int_{t_1}^{t_2} \frac{\sqrt{\alpha^2(t) + [\alpha'(t)]^2}}{\alpha^2(t)} dt$$

where t is the angle between OP_t and horizontal line.

Proof. Let $OP_t = \alpha(t)$, $OP'_t = \beta(t)$ and let $t_1 < t_2$.

$$\text{Then } \int_{t_1}^{t_2} \left| \frac{d\beta(t)}{dt} \right| dt = \int_{t_1}^{t_2} \sqrt{\beta^2(t) + \left[\frac{d\beta(t)}{dt} \right]^2} dt.$$

From (2.1.a), we have

$$\begin{aligned}
 & \int_{t_1}^{t_2} \sqrt{\beta^2(t) + \left[\frac{d\beta(t)}{dt} \right]^2} dt \\
 &= \int_{t_1}^{t_2} \sqrt{\left[\frac{R^2}{\alpha(t)} \right]^2 + \left[\frac{d}{dt} \frac{R^2}{\alpha(t)} \right]^2} dt \\
 &= R^2 \int_{t_1}^{t_2} \sqrt{\left(\frac{1}{\alpha(t)} \right)^2 + \left[-\frac{1}{\alpha^2(t)} \frac{d\alpha(t)}{dt} \right]^2} dt \\
 &= R^2 \int_{t_1}^{t_2} \frac{\sqrt{\alpha^2(t) + [\alpha'(t)]^2}}{\alpha^2(t)} dt.
 \end{aligned}$$

Thus we have the result.

Example 3.3 Let the circle through center of inversion circle $(O)_R$

be $\alpha(t) = \cos t$ and let $0 \leq t \leq \frac{\pi}{3}$.

Then we have

$$\begin{aligned}
 \int_0^{\frac{\pi}{3}} \left| \frac{d\beta(t)}{dt} \right| dt &= R^2 \int_0^{\frac{\pi}{3}} \frac{\sqrt{\cos^2 t + \sin^2 t}}{\cos^2 t} dt \\
 &= R^2 \int_0^{\frac{\pi}{3}} \sec^2 t dt \\
 &= R^2 [\tan t]_0^{\frac{\pi}{3}} \\
 &= \sqrt{3}R^2.
 \end{aligned}$$

On the other hand, in virtue of (2.1.a), if $t = \frac{\pi}{3}$,

$$\beta(t) = \frac{R^2}{\alpha(t)} = \frac{R^2}{\cos t} = 2R^2.$$

Thus $PQ = \sqrt{3}R^2$ (Fig 2.1).



REFERENCES

- [1] Richard S. Millman and George D. Parker (1977), *Elements of Differential Geometry*, Prentice-Hall.
- [2] Barrett O'Neill, *Elementary Differential Geometry*, Academic press.
- [3] Mandredo P. Do Carmo (1976), *Differential Geometry of Curves and Surfaces*, Prentice-Hall, Inc.
- [4] Claire Fisher Adler (1967), *Modern Geometry*, McGraw-Hill, Inc.
- [5] Marvin Jay Greenberg (1974), *Euclidean and Non-Euclidean Geometries*, W. H. Freeman and Company.

< 조 록 >

Inversion의 한 곡선의 길이

양창흥

제주대학교 교육대학원 수학교육전공

지도교수 현진오

중심이 O 이고 반지름의 길이가 R 인 원 $(O)_R$ 에서 두 점 P, P' 가 중심 O 의 같은 쪽에 있고, $OP \cdot OP' = R^2$ 을 만족할 때, 이 두 점 P, P' 을 서로 역(inverse)이라 하고, $(O)_R$ 를 전위 원(inversion circle)이라고 하며, 점 P 에서 P' 으로 보내어 주는 변환을 전위(inversion)라고 한다.

이 논문에서는 2차 Euclid 공간의 곡선으로 제한하여, 전위(inversion)에 의한 $(O)_R$ 의 내부의 곡선 $\alpha(t)$ 에 대응하는 새로운 곡선 $\beta(t)$ 의 길이는 곡선 $\alpha(t)$ 의 길이의 스칼라배로 나타낼 수 있음을 보였다.

* 본 논문은 1993년 8월 제주대학교 교육대학원 위원회에 제출된 교육학 석사학위 논문임.